

National Drought Management Authority

NAROK COUNTY

DROUGHT EARLY WARNING BULLETIN FOR MAY 2018



A Vision 2030 Flagship Project



MAY EW PHASE

Drought Situation & EW Phase Classification

Biophysical Indicators

- **Rainfall**
- There was an early onset of long rains, which started in third dekad of February. The county recorded above normal rains during the month under review. The rains subsided during the third dekad of May.
- **Vegetation condition:**
- The vegetation greenness as depicted by the vegetation condition index (VCI) is normal across all the sub counties.

Socio Economic Indicators (Impact Indicators)

Production indicators:

- The on-going farming activities include weeding Maize and beans in Narok North, Narok East, Narok South, Transmara East and Kilgoris sub-counties. Harvesting of green maize in Narok North, Narok South and Transmara region is also taking place.. Most crops are at flowering and grain filling stage.
- Livestock body condition ranged from fair to good in the pastoral livelihood zone and good in the mixed farming and the agro-pastoral livelihood zones.
- Milk production is stable and within the normal range compared to the long term average.

Access indicators:

- Terms of trade were not favorable in the mixed farming and the agro-pastoral livelihood zones.
- Milk consumption ranged from fair to good in all the livelihood zones.
- Water access for both human and livestock was good in all the livelihood zones

Utilization Indicators:

- Malnutrition cases have gone down and are below the normal range as indicated by percent of mid upper arm Circumference (MUAC).

Early Warning (EW) Phase Classification

| Livelihood Zone | Phase | Trend |
|--|---------------|--------------------|
| AGRO PASTORAL | NORMAL | STABLE |
| MIXED FARMING) | NORMAL | STABLE |
| PASTORAL | NORMAL | STABLE |
| COUNTY | NORMAL | STABLE |
| Biophysical Indicators | Value | Normal Range/Value |
| Rainfall (% of Normal) | 164% | >90% |
| VCI-3Month | 54.75 | >35 |
| Forage condition | Good | Good |
| Production indicators | Value | Normal |
| Crop Condition (Maize) | Normal | Normal |
| Livestock Body Condition | Fair-good | Good |
| Milk Production in litres | 3.5 | >5 |
| Livestock Migration Pattern | Normal | Normal |
| Livestock deaths (Drought related)) | No death | No death |
| Access Indicators | Value | Normal |
| Terms of Trade (ToT) | 74.5 | >100 |
| Milk Consumption in litres | 2.5 | >4 |
| Return distance to water sources in km | 2.2 | <5 |
| Cost of water at source in Kshs (20 litres) | 2-5 | <5 |
| Utilization indicators | Value | Normal |
| Nutrition Status, MUAC (% at risk of malnutrition) | 6 | <10 |

| | | | |
|--|---|---|---|
| <ul style="list-style-type: none"> ▪ Short rains harvests ▪ Short dry spell ▪ Reduced milk yields ▪ Increased HH Food Stocks ▪ Land preparation | <ul style="list-style-type: none"> ▪ Planting/Weeding ▪ Long rains ▪ High Calving Rate ▪ Milk Yields Increase | <ul style="list-style-type: none"> ▪ Long rains harvests ▪ A long dry spell ▪ Land preparation ▪ Increased HH Food Stocks ▪ Kidding (Sept) | <ul style="list-style-type: none"> ▪ Short rains ▪ Planting/weeding |
| | | | |
| Jan | Feb | Mar | Apr |
| May | Jun | Jul | Aug |
| Sept | Oct | Nov | Dec |

1. CLIMATIC CONDITIONS

1.1 RAINFALL PERFORMANCE

- There was an early onset of long rains which started in the third dekad of February. The rains have continued throughout the months of March, April and mid-May. The rains have ceased towards the end of May and the cold spell has kicked off.
- There were light showers in May which were evenly distributed in both time and space in all parts of the county.

1.2 AMOUNT OF RAINFALL AND SPATIAL DISTRIBUTION




Figure 1

- The above satellite image highlights the general performance of the rains in the county for the month of May. There has been a decline in the amount of rainfall received in May compared to last month as shown in Figure 1 above.

2. IMPACTS ON VEGETATION AND WATER

2.1 VEGETATION CONDITION




Figure 2

The vegetation greenness as depicted by the vegetation condition index (VCI) shows normal vegetation greenness as shown in Figure 2.

2.1.1 Field observations

2.1.1.1 Pasture

- The pasture quantity and quality in mixed farming and agro pastoral livelihood zones remained good. In pastoral livelihood zone, pasture conditions ranged from fair to good. The pasture conditions in Mosiro Ward in Narok East sub county and Naroosura ward in Narok south sub county have improved compared to last month due to rains received in the county.
- The available pasture is likely to last for more than four months in mixed farming and agro-pastoral livelihood zones. In pastoral livelihood zone, pastures are likely to last for more than three months.
- The current pasture situation is within the normal range.

2.1.1.2 Browse

- The quality and quantity of browse in all the livelihood zones was good. The available browse is expected to last for more than six months in mixed farming livelihood zone while in pastoral livelihood zone, it is likely to last for more than four months.
- The current browse situation is above the normal range.

2.2 Water resources

2.2.1 Sources

- The main sources of water for livestock and human consumption in the county were boreholes, pans/dams, springs, rivers, traditional river wells and shallow wells as shown by Figure 3.
- The quantity and quality of water was generally good in all the livelihood zones with pans and dams being the most used.
- In pastoral livelihood zone, water quality ranged from fair to good even though some of the pans are silted. In mixed farming and agro pastoral livelihood zones, water quality was good and this was attributed to recharge of water sources in the county.
- The current water sources are expected to last for more than five months in mixed farming and agro-pastoral livelihood zones. In pastoral livelihood zone, the water is likely to last for four months.




Figure 3

- The status of main sources of water is above normal at this time of the year.

2.2.2 Household access and Utilization




Figure 4

The current trekking distances to water points were below the normal range compared to the long term means as shown by Figure 4.

2.2.3 Livestock access

The average distance to main water sources from grazing areas reduced by 35% compared to the previous month. Pastoral livelihood zone recorded the longest distance of five km while the shortest distance was recorded in the mixed farming




Figure 5

livelihood zone at 0.3 km. The current average trekking distance is below the normal range at this time of the year as shown in Figure 5.

The average distance to watering points for households remained stable compared to the previous month.

Pastoral livelihood zone recorded the longest average return distance of three km having decreased from four km last month while the shortest average distance was recorded in mixed farming livelihood zone. The reduction in trekking distance is due to ongoing rains and most household can now access water within the vicinity of their homesteads.

3.0 PRODUCTION INDICATORS

3.1 LIVESTOCK PRODUCTION


3.1.1 Livestock Body Condition

- The cattle body condition was generally good in the mixed farming and some parts in the agro-pastoral livelihood zones. Cattle body condition ranged from fair to good in the pastoral livelihood zone. The body condition for sheep and goats was good across all livelihood zones.
- The stability in livestock body condition is attributed to improved forage condition and water availability due to ongoing rains in the county.
- The current livestock body condition is normal at this time of the year.

3.1.2 Livestock Diseases

- There were no alarming disease outbreaks in the county except cases of blue tongue and pneumonia in small stocks which were being reported in Narok South, Narok East and Narok West sub-counties. This is due to heavy rains and chilly weather conditions being experienced in the county

3.1.3 Milk Production



The average milk production per household per day remained stable compared to the previous month. The stability in milk production was attributed to improved livestock body condition as well as pasture and water availability in the county.

The current milk production rate is below the normal range at this time of the year as shown by Figure 6.

Figure 6

3.2 RAIN-FED CROP PRODUCTION

3.2.1 Stage and Condition of food Crops

- The crops planted in the County include beans, sorghum, maize, wheat and potatoes in Narok North, Kilgoris and Emurua Dikirr sub-counties.
- Minimal weeding of maize and beans in ongoing in some areas.
- The performance of crops was generally good and high yield is expected this season.
- However, some crops have been damaged due to excess rains and floods. In Mosiro ward, some crops within the irrigation scheme were swept away. In Transmara East and West sub counties, crops that were planted earlier have been damaged due to water logging.

4.0 MARKET PERFORMANCE

4.1 LIVESTOCK MARKETING

4.1.1 Cattle Prices




Figure 7

The average price for the medium sized cattle slightly increased by nine percent compared to the previous month.

The highest prices were recorded in the pastoral livelihood zone at Kshs 22,000 per head and the lowest in the agro-pastoral livelihood zone at an average price of Kshs 16,500 per head.

The current cattle prices are above the normal range compared to long term means at this time of the year as shown by Figure 7.

4.1.2 Goat Prices

The average goat price slightly increased by five percent compared to the previous month. The price increase is attributed to good body condition and forage availability in all the livelihood zones. The highest price was recorded in the agro-pastoral livelihood zone at Kshs 4,500 while the lowest price was recorded in the pastoral livelihood zone at Kshs 3,400. The




Figure 8

average goat price was above the normal range at this time of the year as shown by Figure 8.

4.2 CROP PRICES

4.2.1 Maize

The average price of maize per kilogram reduced by 13 percent compared to the previous





Figure 9

month. The highest prices were recorded in pastoral livelihood zone at Kshs 74 per kg while the lowest was recorded in the mixed farming livelihood zone at an average price of Kshs 37 per kg. The overall price is below the normal range as shown by Figure 9.

4.2.2 Beans

- The average price of beans increased by seven percent compared to the previous month. The highest price was recorded in the agro-pastoral livelihood zone at Kshs. 123/kg compared to the mixed farming livelihood zone which recorded a lower price of Kshs.90 per kg.
- The current price is below the normal range at this time of the year as shown



by Figure 10.

Figure 10

4.3 Livestock Price Ratio/Terms of Trade




Figure 11

The cereal/meat price ratio is above the normal range at this time of the year as shown in Figure 11.

The Terms of Trade (ToTs) based on the Cereal/Meat price ratio increased by 12 percent compared to last month. The increase was attributed to decrease in maize price and increase in goat price. This implies that a sale of one goat fetched 74.5kg of cereals. The ToTs were less favourable in the agro-pastoral and mixed farming livelihood zones compared to pastoral livelihood zone.

5.0 FOOD CONSUMPTION AND NUTRITION STATUS

5.1 MILK CONSUMPTION




Figure 12


consumption rate at 1.8 litres. The current milk consumption rate is within normal range at this time of the year as shown by Figure 12.

Average household milk consumption per day slightly increased compared to the previous month. There was significant disparity in milk consumption across the livelihoods with agro pastoral livelihood zone having the highest consumption rate at five litres, followed by mixed farming zone at six litres while

the pastoral zone had the least

5.2 FOOD CONSUMPTION SCORE


- About 67%, 17% and 16% of the households fell within the acceptable, borderline and poor food consumption score categories respectively in the month under review.
- The poor food consumption score implies household are not consuming a balanced diet and rarely consuming food rich in protein. Borderline implies that households are consuming staples and vegetables every day accompanied by oil and F



- However, there was a significant variation across livelihood zones as shown in Figure 13 with mixed farming and pastoral livelihood zones having the bulk of their populations with acceptable food consumption scores. In agro-pastoral livelihood zone, the majority of the population had poor food consumption scores. The inter-livelihood variation is attributed to high consumption of meat and milk in most households in the pastoral and mixed farming livelihood zones compared to those in the agro-pastoral livelihood zone.

5.3 HEALTH AND NUTRITION STATUS

5.3.1 Nutrition Status



The percentage of children less than five years of age who are at risk of malnutrition as indicated by the Mid-Upper Arm Circumference (MUAC) reduced compared to the previous month. The pastoral and the agro-pastoral livelihood zones recorded the highest number of children at risk of malnutrition at six percent compared to mixed farming livelihood zone which recorded one percent. The highest number of children at risk of malnutrition was still being witnessed in Mosiro and Naroosura

Figure 14

wards in pastoral livelihood zone owing to poor dietary diversity coupled with low household milk consumption and high food prices. The dietary diversity in mixed farming livelihood zone has improved slightly due to availability of green vegetables and cereals. This zone harvested maize in January and February which contributed positively to children's nutrition status coupled with improved milk intake. The current rate of malnutrition has improved and is below the normal range compared to the long term average as shown by Figure 14.

5.3.2 Health

- The county experienced increased cases of fever due to change of weather though it is not alarming.

5.4 COPING STRATEGIES




Figure 15

Households employed less unusual coping strategies in accessing food across all the livelihood zones as shown by Figure 15. The pastoral livelihood zone had higher CSI compared to other zones. This implies that households in the pastoral livelihood zone employed more coping strategies compared to those in other zones who can sell livestock and livestock products to purchase food. The coping strategies are normal at this time of the year

6.0 CURRENT INTERVENTION MEASURES (ACTIONS)

6.1 NON-FOOD INTERVENTIONS

- Malnutrition screening has been concluded. It was being conducted by NDMA in collaboration with Ministry of Health through EU/DCF support in Narok East, Narok South and Transmara East Sub counties.
- The County Government in collaboration with the National Government and the World Bank have started implementing NARIGP project. (National Agricultural Rural Investment Growth Project) targeting small holder farms in 20 Wards of Narok County.
- The Regional Pastoral Livelihood Resilience Project is undertaking some drought response interventions which include livestock restocking, animal feeds supplementation of 6,000 bags of range cubes and 5,000 pieces of mineral blocks. Beneficiary targeting has been done and distribution logistics are being worked on.

6.2 FOOD AID

- No food aid was issued during the month.

7.0 EMERGING ISSUES

7.1 Insecurity/Conflict/Human Displacement

- No major conflict issues were observed during the month under review.

7.2 Migration

- There were no incidences of migration in the county.

7.3 FOOD SECURITY PROGNOSIS

- The performance of the long rains season was above normal. This has had a positive impact on various food security sectors. Forage conditions have significantly improved, a factor which is likely to enhance livestock productivity for the next three months. Water situation is expected to stabilise for the next two months. Malnutrition cases are likely to go down further due to expected improvement in milk production and consumption. However, there is likely to be a delayed harvest of some crops which were affected by excessive rains that led to water logging which hampered planting and germination of crops.

8.0 RECOMMENDATIONS

- Some limited vaccination against blue tongue disease is required in Narok West and Narok South sub counties.
- Water hygiene and sanitation standards must be observed in order to minimise chances of outbreak of water borne diseases.
- Repair of damaged infrastructure due to heavy rains particularly rural access roads. The Ntulele-Mosiro road should be given top priority to ease movements of goods to the markets.