


**National Drought Management Authority  
LAIKIPIA COUNTY  
DROUGHT EARLY WARNING BULLETIN FOR JUNE 2018**



A Vision 2030 Flagship Project



**JUNE 2018 EW PHASE: NORMAL**



**Drought Situation & EW Phase Classification**

**Biophysical Indicators**

**Rainfall:**

- In the month of June, the County experienced 6-13 days of moderate to heavy rainfall. The rainfall was generally well distributed in terms of time and space.

**Vegetation Condition:**

- The Vegetation Condition Index (VCI) is above the normal range for the period, indicating very good state of pasture and browse condition across most areas.
- The available pasture and browse is expected to last for at the least 3 months.

**Socio Economic Indicators (Impact Indicators)**

**Production Indicators:**

- There was no migration recorded.
- Milk production per household was within the normal range at this time of the year.
- The body condition of animals was within the normal range for the period.

**Access indicators:**

- The terms of trade are above the normal range
- Milk consumption is within the normal range
- The return distance from water sources to grazing areas is also within the normal range.

**Utilization indicators:**

- All within the normal range.

LIVELIHOOD ZONE	EW PHASE	TREND
PASTORAL	Normal	Stable
MMF	Normal	Stable
MF	Normal	Stable
<b>COUNTY</b>	<b>Normal</b>	<b>Stable</b>
Biophysical Indicators	Value	Normal range
% of Average rainfall	258%	80-120%
SPI-3 month (TAMSAT)	-	-1 to 1
VCI (1 month)	84.4	35-50
State of Water Sources	5	5
Production indicators	Value	Normal range
Livestock Migration Pattern	No Migration	No Migration
Livestock Body Condition	5	4-5
Milk Production (Lt)	5.7	4.9
Reported livestock deaths	No death	No death
Crops area planted (%)	-	% of LTA
Access Indicators	Value	Normal ranges
Terms of Trade (ToT)	113.9	90
Milk Consumption (Lt)	2	>1.7
Return Distance (Water Sources - grazing areas)	3	< 5
Return Distance water to Grazing areas (Pastoral)	3.5	<7
Utilisation indicators	Value	Normal ranges
MUAC (Mid at risk)	1.8	< 18
Coping Strategy Index (CSI)	-	<1


<ul style="list-style-type: none"> <li>▪ Short rains harvests</li> <li>▪ Short dry spell</li> <li>▪ Reduced milk yields</li> <li>▪ Increased HH Food Stocks</li> <li>▪ Land preparation</li> </ul>	<ul style="list-style-type: none"> <li>▪ Planting/Weeding</li> <li>▪ Long rains</li> <li>▪ High Calving Rate</li> <li>▪ Milk Yields Increase</li> </ul>	<ul style="list-style-type: none"> <li>▪ Long rains harvests</li> <li>▪ A long dry spell</li> <li>▪ Land preparation</li> <li>▪ Increased HH Food Stocks</li> <li>▪ Kidding (Sept)</li> </ul>	<ul style="list-style-type: none"> <li>▪ Short rains</li> <li>▪ Planting/weeding</li> </ul>								
Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sept	Oct	Nov	Dec

# 1 CLIMATIC CONDITIONS

## 1.1 Rainfall Performance


### Performance

- During the month under review (June), Laikipia County recorded 6 to 13 days of heavy to moderate rainfall. The marginal mixed farming (MMF) livelihood zones recorded 6 days of moderate to light showers with good distribution in terms of space. The mixed farming (MF) livelihood zone recorded 13 days of heavy rainfall. The Pastoral livelihood zone recorded 2 days of moderate to light showers with fair distribution.



- The rains received in June were approximately 258% of the long-term average expected amount for the month, which is way above normal. This is higher compared to 205% recorded in May.

## 1.2 Amount of Rainfall and Spatial Distribution




- According to the chart above, the amount of rain received in June amounted to 133.2 mm for the month, which is above the long term average of 46.7 mm by the same time hence way above the normal range.
- Compared to May at 126.5 mm, the precipitation levels increased slightly. This is attributed to the persistent rains even as the season moves towards cessation.
- The rainfall distribution was very good in terms of time and space across all the livelihood zones.

## 2. IMPACT ON VEGETATION AND WATER

### 2.1 Vegetation Condition


#### 2.1.1 Vegetation Condition Index (VCI)



- The VCI matrix above indicates an above normal vegetation greenness for the month of June, which is the same compared to the previous month. From field observations, this was also the case as vegetation thrived in both quality and quantity.
- The state of vegetation is attributed to the effect of the above normal rains. All wards reported above normal vegetation condition for the month.
- The actual VCI (three month) at 84.4 is well above the long-term average for the month, a tremendous improvement compared to 69.87 in May.

#### 2.1.2 Pasture


- Key informant interviews indicated that the pasture condition was overall good (100%) as shown in the chart below.



- Compared to the previous month (95.8% good and 4.2% fair), there was a substantial improvement of the pasture condition in both quantity and quality.
- The improvement in pasture condition is attributed to the positive effect of the above normal long rains (MAM).
- There was no major constraint to pasture access.
- The quantity of pasture available is expected to last 3 months in the Pastoral and the MMF zones. In the MF zone the pasture condition is expected to last for at least 3-4 months.

### 2.1.3 Browse

- According to the key informants interviewed, the browse condition was split between good (87.5%) and fair (12.5%) as shown in the chart below.




- Compared to the previous month (good – 79.2%, fair – 20.8%), the browse condition recorded a significant improvement.
- The improved state of the browse condition is attributed to the above normal long rains (MAM).
- No major constraint to browse access was recorded.
- The quantity of browse available is expected to last 3-4 months in the Pastoral and the MMF zone. In the MF zone the pasture condition is expected to last for 4 months.

## 2.2 Water Resource

### 2.2.1 Sources


- During the month under review, the main water sources for domestic and livestock use in the County were pans and dams (28.6%), boreholes (25.8%), shallow wells (23.2%) and rivers (14.3%). Others were traditional river wells (7.1%).



- Compared to the previous month, the quantity and quality of water has remained stable due to the prevailing rains. The main water source has shifted from boreholes to pans and dams.
- The main water sources are expected to last as follows: - Pastoral (boreholes - permanent, pans and dams – 2-3 months), MMF (borehole - permanent, rivers – 3 months, pans and dams – 2-3 months), MF (shallow wells - 3 months, traditional river wells – 4 months, pans and dams – 3 months).

## 2.2.2 Household Access and Utilization


- In June, the average return distances from households to water sources remained low at 2.6 km, slightly higher than the previous month. The furthest return distance of 2.7 km was recorded in the marginal mixed farming zone (mainly Laikipia East and parts of West) slightly up from 2.5 km in May.



- The low distances are attributed to the positive effect of the MAM rains. The slight increase in distances is attributed to the changing water sources as the MAM rains cease.

## 2.2.3 Livestock Access

- The average return distance from water sources to grazing areas slightly increased to 3 km, up from 2.9 km in May, which is an insignificant increase. The longest return distance of 3.5 km was recorded in the Pastoral zones, a slight increase compared to 3.4 km in May. MMF zones recorded 2.2 km, a slight decrease from 2.2 km the previous month.



- The low distances from water sources to grazing areas can be mainly attributed to the low distances covered in search of water and pasture.

### **3 PRODUCTION INDICATORS**

#### **3.1 Livestock Production**

##### **3.1.1 Livestock Body Condition**

- Across the county, the livestock body condition was classified at level 5 (normal – good smooth appearance).
- The livestock body condition was good for cattle and shoats. This is normal at this time of the year.
- Compared to same time last year, the body condition of livestock is much better.
- The livestock body condition has improved compared to last month due to the low trekking distance in search of pasture and water coupled with the increase of pasture quantity and quality.
- The body condition of cattle and shoats was good across all the livelihood zones.

##### **3.1.2 Livestock Diseases and Deaths**

- There were few unconfirmed cases of FMD reported in the Pastoral Zones.

##### **3.1.3 Milk Production**

- Across the county, the sampled households recorded an increased average milk production of 5.7 litres per household per day in June, up from 5.5 litres in May. The Pastoral zone recorded the least milk production per household at 2.2 litres, a 9% increase compared to the previous month. This milk was obtained from cattle.
- The increase in milk production is attributed to the low trekking distance in search of pasture and water and the increase of pasture quantity and quality.
- The milk production is slightly above the average levels (4.9 litres per household) at this time of the year.

#### **3.2 Rain-fed Crop Production**


##### **3.2.1 Stage and Condition of Food Crops**

- Currently, crops that are found in most farms in the MF and MMF zones are maize, wheat, potatoes and beans. The crops are doing well although threat of armyworm is of a great concern to farmers. Maize is at flowering stage, beans at podding stage whereas potatoes are at flowering and second moulding stages. Wheat is at piping stage. Most farmers are applying second fertiliser (Top dressing) and second weeding. The current excessive rains have resulted in leaching in farms in the crop growing zones.
- Cases of armyworm and other pests have been reported in farms in the MMF zones, particularly Olmorani, Sosian and Ngobit wards.

## 4 MARKET PERFORMANCE


### 4.1 Livestock Marketing

#### 4.1.1 Cattle Prices (Market)



- In June, the county recorded an average cattle price of Kshs. 32,041 at the markets, a significant increase (by 6%) compared to the previous month. This can be attributed to the good body condition as a result of low trekking distances in search of pasture and water.
- The pastoral zone recorded the highest cattle price whereas MMF recorded the lowest of the same.
- Compared to the long term average, the current price is way higher than the long term average hence above normal.


#### 4.1.2 Small Ruminants Prices (Goat)



- During the month under review, the average price of a goat in Laikipia was recorded at Kshs. 3,917, which is a 2% increase compared to the previous month. The slight increase can be attributed to the good body condition of the animals at the markets.
- The highest average goat price was recorded in the MF zone whereas the lowest price was recorded in the MMF zone. The current goat price is slightly higher by 13.5% compared to the long term average, hence above the normal range.


## 4.2 Crop Prices

### 4.2.1 Maize (market price)



- During the month under review, an average maize price of Kshs. 34 was recorded at the markets as shown in the graph above, same as the previous month. The low market price is attributed to the availability of maize at household level because of the ongoing piecemeal harvests, previous maize harvests and robust supply from other counties.
- The highest average market price of maize at Kshs.40 was recorded in Kimanjo (Pastoral) and Sirima (MMF) markets whereas the lowest at Kshs. 25 were recorded in Olmoran (MMF) market.
- Compared to the three-year average, the current price is lower by 13 shillings.

### 4.3 Livestock Price Ratio/ Terms of Trade



- From the graph above, the June average price of a goat at Kshs. 3,917 was able to purchase 113.93 Kg of maize, slightly higher than the previous month.
- The current ToT (Terms of Trade) can be attributed to the robust goat prices and low maize prices across all livelihood zones. The ToT favours livestock keepers as they are still able to purchase more maize for the price of a goat.
- When compared to the three-year average, the ToT is above normal.



#### **4.4 Implication on Food Security**


- The stable cereal prices have ensured that households have access to food hence a stable food secure status across the county.
- The good body condition resulting from adequate water and pastures has ensured stable supply of milk and robust prices at the markets. This has had a positive effect on both household food security and purchasing power.
- The almost ceasing enhanced MAM rains have exceeded expectation and have had an overall positive effect on forage, crops and water sources. As a result, the stability of food prices and availability of water and pasture for livestock is ensured for the near future.

## 5 FOOD CONSUMPTION AND NUTRITION STATUS

### 5.1 Milk Consumption

- During the month under review, the sampled households recorded an average milk consumption of 2 litres per day, a slight increase compared to the previous month (at 1.9), with most of the milk coming from cattle.
- The milk consumption levels are within the normal levels (>1.8 litres) expected at this time of the year.
- For the MMF and MF zones, the larger percentage of the milk produced (75% and 58% respectively) was sold as households sought to raise income for other household needs whereas for Pastoral zones, 99% of the milk produced was used to supplement the diet.


### 5.2 Food Consumption Score



- According to the chart above, all the households in the mixed farming livelihood zone maintained an acceptable food score. The Pastoral zone followed with an acceptable food score of 90%, a borderline food score of 8.3% and a poor food score of 1.7%.
- 69.3% of the households in the marginal mixed farming (MMF) zone had an acceptable score whereas 29.5% had a borderline score.
- In general, there was a slight increase in dietary diversity in all the livelihood zones.

## 5.3 Health and Nutrition Status

### 5.3.1 Nutrition Status



- The percentage of children under-five years of age who are at risk of malnutrition is 1.8%, slightly higher than the previous month, which is an insignificant difference.
- There were no reported cases falling under SAM and MAM for the current month.

### 5.3.2 Health

- There were no reported major diseases apart from cases of common cold affecting both adults and children across the sentinel sites during the period under review in the sampled households.

## 5.4 Consumption based coping strategies

- The most common types of the strategies employed were relying on less preferred food, taking fewer meals, borrowing and purchasing food on credit.

## 6 CURRENT INTERVENTION MEASURES (ACTION)

### 6.1 Non-Food Interventions

- No non – food interventions were reported in the County during the period.

### 6.2 Food Aid

- The community around Survey (MMF) received food aid from the County Government of Laikipia.

## 7 EMERGING ISSUES

### 7.1 Insecurity/Conflict/Human Displacement

- Case of livestock theft(Shoats) has been reported at Ilgwesi (Pastoral zone)
- Cases of water logging in farms has been reported at Kinamba (MF), destroying crops in the farms.
- Traders have had challenges in accessing nearby markets in Tigithi, Salama, Ngobit and Mukogodo wards due to poor roads in these zones as a result of heavy downpour observed previously.

### 7.2 Migration

- During the month under review, no cases of both human and animal migration were reported in the County.

### 7.3 Food Security Prognosis

- The long rains have performed beyond expectation and this has led to the tremendous improvement of the forage condition and water. As the rains cease, their positive effect across all indicators, i.e. the resultant increase in pasture and browse condition has resulted in the improvement of the livestock body condition, leading to better prices at the markets and stable food prices.
- Generally, a good crop yield is expected except in areas where leaching and flooding has occurred or armyworms crop pests have been reported. The county is expected to remain food secure for the next three to four months.

## 8 RECOMMENDATIONS

- Enhance crop pests and disease surveillance, foster extension services on the same **Action: County Govt. – agriculture.**
- Carry out water borne disease surveillance, advice communities on hygiene **Action: County Govt (health and water).**
- Initiate long-term interventions geared towards curbing human wildlife conflicts. **Action: KWS.**
- Provision of certified fertilizers at subsidized rate to farmers **Action: FAO, ASDSP, County Govt, National Govt, and Private Stakeholders.**
- Capacity building farmers on Conservation Agriculture as a modern method of farming **Action: FAO, ASDSP, County Govt, and Private Stakeholders.**
- Continue drought drought contingency plans review and start preparedness projects **Action: NDMA.**

## REFERENCE TABLES

**Table 1: Drought Phase Classification**

Normal	Alert	Alarm	Emergency
All environmental Agricultural and pastoral indicators are within the seasonal ranges	Biophysical drought indicators move outside seasonal ranges	Environmental and at least three production indicators are outside long term seasonal ranges	All Environmental, Metrological and Production indicators are outside normal ranges.
<b>Recovery:</b> The drought phase must have reached at least Alarm stage. Recovery starts after the end of drought as signalled by the environmental indicators returning to seasonal norms; local economies starting to recover			

**Table 2: Standardized Precipitation Index (SPI)**

Color	SPI Values	Meteorological Drought Category
	> +1.5 or more	Wet Conditions
	0 to +1.5	No drought
	-0.1 to -0.99	Mild drought
	-1 to -1.99	Severe drought
	<-2 and less	Extreme drought

**Table 3: Vegetation Condition Index Values (VCI)**

Color	VCI values	Agricultural Drought Category
	<b>3-monthly average</b>	
	≥50	Wet
	35 to 50	No agricultural drought
	21 to 34	Moderate agricultural drought
	10 to 20	Severe agricultural drought
	<10	Extreme agricultural drought

**Table 4: Livestock Body Condition**

Level	Classification	Characteristics (this describes majority of the herd and not individual isolated Stock)
5	Normal	Very Fat Tail buried and in fat
		Fat, Blocky. Bone over back not visible
		Very Good Smooth with fat over back and tail head
		Good smooth appearance
4	Moderate	Moderate. neither fat nor thin
3	Stressed	Borderline fore-ribs not visible. 12th & 13th ribs visible
2	Critical	Thin fore ribs visible
1	Emaciated	Very thin no fat, bones visible
		Emaciated, little muscle left

### Definition of Early Warning Phases

The EW phases are defined as follow:

**NORMAL:** The normal phase occurs when **biophysical drought indicators (VCI and SPI) show no unusual fluctuations** hence remain within the expected ranges for the time of the year in a given livelihood zone, division or county

**ALERT:** The alert phase is when either the **vegetation condition index or the standard precipitation index (biophysical indicators) show unusual fluctuations below expected seasonal ranges** within the whole county/sub-county or livelihood zones.

**ALARM:** The alarm phase occurs when both **biophysical and at least three production indicators fluctuate outside expected seasonal ranges** affecting the local economy. The production indicators to be considered are livestock body condition, crop condition, milk production, livestock migration and livestock mortality rate.

If **access indicators** (impact on market, access to food and water) move outside the normal range, the status remains at “alarm” but with a worsening trend. Proposed access indicators include ToT, price of cereals, availability of cereals and legumes, and milk consumption. The trend will be further worsening when also welfare indicators (MUAC and CSI) start moving outside the normal ranges.

**EMERGENCY:** In the emergency phase, **all indicators are outside of normal ranges**, local production systems have collapsed within the dominant economy. The emergency phase affects asset status and purchasing power to extent that seriously threatens food security. As a result, coping strategy index, malnutrition (MUAC) and livestock mortality rates move above emergency thresholds.

**RECOVERY:** **Environmental indicators returning to seasonal norms.** The drought phase must have reached at least Alarm stage. Recovery starts after the end of drought as signalled by the environmental indicators returning to seasonal norms while production indicators are still outside the normal seasonal range but local economies start to recover. The status changes to normal once the bio physical and production indicators are back to normal range.